

SOLUTIONS

PROGRESS TEST-6

RBA

JEE MAIN PATTERN

Test Date: 04-12-2017



**Corporate Office: Paruslok, Boring Road Crossing, Patna-01
Kankarbagh Office: A-10, 1st Floor, Patrakar Nagar, Patna-20
Bazar Samiti Office : Rainbow Tower, Sai Complex, Rampur Rd.,
Bazar Samiti Patna-06
Call : 9569668800 | 7544015993/4/6/7**

PHYSICS

1. (B) Component of \vec{v} along $\hat{a} = (\vec{v} \cdot \hat{a})$

$$= (6\hat{i} + 2\hat{j} - 2\hat{k}) \cdot \frac{(\hat{i} + \hat{j} + \hat{k})}{\sqrt{3}}$$

$$= \frac{6+2-2}{\sqrt{3}} = \frac{6}{\sqrt{3}} = 2\sqrt{3}$$

In vector form = $(2\sqrt{3}) \hat{a}$

$$= 2\sqrt{3} \frac{(\hat{i} + \hat{j} + \hat{k})}{\sqrt{3}}$$

$$= 2(\hat{i} + \hat{j} + \hat{k})$$


2. (C)

Let acceleration of mass m relative to wedge down the plane is a_r . Its absolute acceleration in horizontal direction is $a_r \cos 60^\circ - a$ (towards right). Hence, let N be the normal reaction between the mass and the wedge. Then

$$N \sin \theta = Ma = m(a_r \cos 60^\circ - a)$$

$$\text{or } a_r = \frac{(M+m)a}{m \cos 60^\circ} = \frac{2(M+m)a}{m}$$

3. (C)



$$T \sin \theta = 20$$

$$T \cos \theta = W$$

$$\tan \theta = \frac{20}{W}$$

$$W = 20 \cot \theta$$

$$W = 20 \cot 30^\circ$$

$$W = 20\sqrt{3} = 20 \times 1.732$$

$$\text{Weight } W = 34.64 \text{ N}$$

$$\text{mass 'm'} = \frac{W}{g} = 3.5 \text{ kg.}$$

4. (A)

Since 'M' is at rest the tension in the string

$$= \frac{Mg}{2} \text{ Let acceleration of } m \text{ and } m' \text{ is 'f' one will move downward and other will move upward}$$

$$mg - \frac{Mg}{2} = mf \dots\dots(i)$$


$$\frac{Mg}{2} - m'g = m'f \dots\dots(ii)$$

Solving equation (i) and (ii)

$$\frac{4}{M} = \frac{1}{m} + \frac{1}{m_1}$$

5. When the object is placed at the focus of the lens, the refracted rays will be incident normally on the silvered surface. So, they will retrace their path.

Hence, the image will be formed at the location of the object. In this way, the combination behaves as a concave mirror of radius of curvature (R) = 20 cm



$$\therefore f = \frac{R}{2} = 10 \text{ cm}$$

\therefore (D)

6. At maximum height $v_1 = v \cos \theta$

$$\text{At half of maximum height } v_2 = \sqrt{v^2 \cos^2 \theta + (v_y)^2}$$

$$\text{For } v_{\text{vertical}} \quad v_y^2 = v^2 \sin^2 \theta - 2g \frac{H}{2} = v^2 \sin^2 \theta - g \frac{v^2 \sin^2 \theta}{2g} = \frac{v^2 \sin^2 \theta}{2}$$

$$\frac{v_1}{v_2} = \sqrt{\frac{2}{5}} \Rightarrow \tan \theta = \sqrt{3}, \theta = 60^\circ$$

\therefore (D)

$$7. \quad T = \frac{2u \sin \theta}{g} = \frac{2 \times 10 \times \frac{4}{5}}{10} = \frac{8}{5} \text{ s}$$

$$OB = R = \frac{u^2 \sin 2\theta}{g} = \frac{100 \times 2 \times \frac{4}{5} \times \frac{3}{5}}{10} = \frac{48}{5} \text{ m}$$

$$AB = \frac{8}{5} \times 5 = 8 \text{ m}$$

$$OA = OB + AB = \frac{48}{5} + 8 = 17.6 \text{ m}$$

\therefore (D)

8. $v = 2t^2, r = 100 \text{ m}$

$$a_t = \frac{dv}{dt} = 4t$$



$$a_t (t = 5\text{s}) = 20 \text{ ms}^{-2}$$

$$v (t = 5\text{s}) = 50 \text{ ms}^{-1}$$

$$a_c (t = 5\text{s}) = \frac{v^2}{r} = \frac{50 \times 50}{100} = 25 \text{ ms}^{-2}$$

$$a = \sqrt{a_c^2 + a_t^2} = \sqrt{1025} \approx 32 \text{ ms}^{-1}$$

\therefore (D)



$$9. \quad t_1 = \frac{2u \sin \theta}{g}, t_2 = \frac{2u \cos \theta}{g}$$

$$\therefore t_1 t_2 = \frac{2u^2 \sin 2\theta}{g \times g} = \frac{2}{g} R$$


$$\therefore t_1 t_2 \propto R$$

$$\therefore \quad \text{(C)}$$

$$10. \quad \text{(B)} \quad T = m_2 \sqrt{g^2 + \left(\frac{v_2^2}{r}\right)^2} = \frac{m_1 v_1^2}{r}$$

11. (B)

In time t particle has rotated an angle $\theta = \omega t$. Displacement




$$s = PQ = \sqrt{QR^2 + PR^2}$$

$$= \sqrt{(a \sin \omega t)^2 + (a - a \cos \omega t)^2}$$

$$s = 2a \sin \frac{\omega t}{2}$$

12 (D)

$$N \sin \theta = mg$$



$$N \cos \theta = mv^2/r$$


$$\Rightarrow \tan \theta = rg/v^2$$

$$\Rightarrow \frac{r}{h} = \frac{rg}{v^2}$$

$$\Rightarrow h = \frac{v^2}{g} = 2.5 \text{ cm}$$

13. (D)

$$\omega = 2/\pi \text{ rev/s} = 4 \text{ rad/s}$$



$$T \sin \theta = m \omega^2 r$$

$$\Rightarrow T \sin \theta = m \omega^2 (L \sin \theta)$$

$$\Rightarrow T = m \omega^2 L = m(4)^2 L = 16mL$$

14. (D)

\therefore Cyclist returns at initial point

\therefore Displacement = 0

$$v_{av} = \frac{s}{t} = \frac{R + \frac{\pi R}{2} + R}{(1/6)} = 6R(2 + \pi/2)$$

$$= 21.4 \text{ km}$$

15. (B)

$$W_g + W_{Fr} = \Delta K$$

$$W_g = -W_{Fr}$$

$$W_{Fr} = -mgh$$


$$W_f + W_g + W_f = \Delta k$$

$$W_f = -W_g - W_{Fr}$$

$$(W_f = 2mgh)$$

16. (C)

The work done by man is negative of magnitude of decrease in potential energy of chain.



$$\Delta U = mg \frac{L}{2} - \frac{m}{2} g \frac{L}{4} = 3 mg \frac{L}{8}$$

$$\therefore W = - \frac{3mgL}{8}$$

17. (D)

$$\therefore \vec{F} = - \left[\frac{\Delta U}{\Delta x} \hat{i} + \frac{\Delta U}{\Delta y} \hat{j} \right] = -[-7\hat{i} + 24\hat{j}]$$

$$\therefore F = \sqrt{7^2 + 24^2} = 25 \text{ N} \Rightarrow a = F/m = 25/5 = 5 \text{ m/s}^2$$

$$\therefore v = u + at$$

$$= 0 + 5 \times 2 = 10 \text{ m/s}$$

18. (A)

$$\text{Power} = Av^3d$$

$$= (Av)(v^2d)$$

$$= (\text{volume per sec}) \times (\text{pressure})$$

$$= \frac{4000 \times 10^{-6}}{60} \times (130 \times 10^{-3} \times 13.6 \times 10^3 \times 10)$$


$$= 1.15 \text{ watt}$$

$$19. \quad \text{Total frictional force on } AB = \int_x^L \frac{Mg}{L} Kx \, dx$$

\therefore Heat generated

$$= \int_0^L \frac{MgK}{L} \left[\frac{L^2 - x^2}{2} \right] dx = \frac{KmgL^2}{3}$$

\therefore (B)



20. (B)

$$B_{AB} = \frac{\mu_0 I}{4\pi(d)} (\sin\alpha_1 - \sin\alpha_2)$$


Where d is the \perp distance and α_1 and α_2 are the angles between \perp and ends of wire.

$$B_{AB} = \frac{\mu_0 I}{4\pi\sqrt{2}} [\sin 90^\circ - \sin 45^\circ]$$

$$= \frac{\mu_0 I}{4\pi\sqrt{2}} \left[1 - \frac{1}{\sqrt{2}} \right] = \frac{\mu_0 I(\sqrt{2}-1)\hat{k}}{8\pi}$$

Magnetic field due to BC will be same and in same direction.

$$\therefore B_0 = \frac{2 \times \mu_0 I(\sqrt{2}-1)}{8\pi} \hat{k}$$



21. (C)

$$V_{CD} = \phi \times \text{Balance length}$$

$$9.91 \text{ V} = \left(\frac{E_0}{L} \times \frac{R_w}{R_{\text{Net}}} \right) \times B \cdot \ell$$

$$9.91 \text{ V} = \frac{100}{10} \times \frac{991}{1000} \times B \cdot \ell$$

$$B \cdot \ell = \frac{9.91}{9.91} = 1 \text{ m}$$

22. (A)


$$E_0 = \frac{kq}{1^2} + \frac{kq}{2^2} + \frac{kq}{4^2} + \frac{kq}{8^2} + \dots \dots \dots \infty$$

$$E = kq \left(1 + \frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \dots \dots \infty \right)$$

$$E = kq \left(\frac{q}{1-r} \right) = kq \cdot \frac{1}{1 - \left(\frac{1}{4} \right)} = \frac{kq \cdot 4}{3}$$

$$12 \times 10^3 = \frac{9 \times 10^9 \times 4}{3} \times q \quad \therefore q = 1 \mu\text{C}$$

23. (A)

Initially force on charge q_0 is

$$F = q_0(2E) = 2q_0E$$

If separation between the plates is doubled $E =$ same $\therefore F =$ same

$$24. \quad \frac{1}{C_\infty} = \frac{1}{C_\infty + C} + \frac{2}{C} = \frac{3C + 2C_\infty}{C(C_\infty + C)}$$

$$2C_\infty^2 + 2CC_\infty - C^2 = 0 \Rightarrow C_\infty = C \left(\frac{-1 + \sqrt{3}}{2} \right)$$

 \therefore (C)

25. (B)

Potential at centre = $V_q + V_{\text{induced}}$

$$= V_q + 0$$


$$= V_q = \frac{q}{4\pi\epsilon_0 r}$$

Potential at P = potential at C

$$V_q + V_{\text{induced}} = \frac{q}{4\pi\epsilon_0 r}$$

$$V_{\text{induced}} = \frac{q}{4\pi\epsilon_0 r} - \frac{q}{4\pi\epsilon_0 r_1} = \frac{kq}{r} - \frac{kq}{r_1}$$

26. (D)



∴ Force perpendicular to BC is zero.

27. [C]

The capacitor is charged by a battery of 25 V. Let the magnitude of surface charge density on each plate be σ . Before inserting the dielectric slab, electric field strength between the plates,

$$E = \frac{\sigma}{\epsilon_0} = \frac{V}{d}$$

$$\text{or } E = \frac{\sigma}{\epsilon_0} = \frac{25}{5 \times 10^{-3}} = 5000 \text{ N/C}$$

The capacitor is disconnected from the battery but charge on it will not change so that σ has the same value. When a dielectric slab of thickness 3mm is placed between the plates, the thickness of air between the plates will be $5 - 3 = 2$ mm. Electric field strength in air will have the same value (5000 N/C) but inside the dielectric, it will be $\frac{5000}{K} = \frac{5000}{10}$


$$= 500 \text{ N/C}$$

$$\text{so potential difference} = E_{\text{air}} d_{\text{air}} + E_{\text{med}} d_{\text{med}}$$

$$= 5000 \times (2 \times 10^{-3}) + 500 \times (3 \times 10^{-3})$$

$$= 11.5 \text{ V}$$

28. (B)



Step I :

$$\text{Step II : } V_A - V_B = \frac{Kq}{r} - \left(-\frac{Kq}{r} \right) = \frac{2Kq}{r}$$

$$\text{Step III : } C = \frac{q}{V_A - V_B} = \frac{qr}{2Kq} = 2\pi\epsilon_0 r$$

29. (C)

$$U = \frac{1}{2} CV^2$$

$$U = \frac{1}{2} \left[\frac{\epsilon_0 b(\ell - x)}{d} + \frac{\epsilon_0 bx}{d} K \right] v^2 = \frac{\epsilon_0 bv^2}{2d} (\ell - x + xK)$$

$$U = \frac{\epsilon_0 bv^2}{2d} [\ell + k(K - 1)]$$

30 (D)

$$C_{\text{eq}} = C_1 + C_2 = 2 + 2 = 4\mu\text{F}$$

$$q = C_{\text{eq}} V = 4\mu\text{F} \times 8 = 32\mu\text{q}$$

CHEMISTRY

31. (D)

Equivalent mass of KMnO_4 in acidic, basic and neutral medium are 31.6, 158 and 52.6. The ratio will be 31.6 : 158 : 52.6.

$$\begin{array}{lcl} \text{Acid medium} & : & \text{Basic medium} & : & \text{Neutral medium} \\ E_w = \frac{M_w}{5} = \frac{158}{5} & : & E_w = \frac{M_w}{1} = \frac{158}{1} = 158 & : & E_w = \frac{M_w}{3} = \frac{158}{3} \\ 3 & : & 15 & : & 5 \end{array}$$

32. (C)

$$W = \Delta E \quad (\text{For } q = 0)$$

$$-P_{\text{ext}}\Delta V = nC_v\Delta T$$

$$-1(2-1) = 1 \times \frac{3}{2}R(T_f - T)$$

$$\frac{-2}{3R} = T_f - T$$

$$T_f = T - \frac{2}{3 \times 0.0821}$$

33. (C)

The van der Waals' equation for 1 mol gas will be

$$\left[P + \frac{a}{V^2} \right] [V - b] = RT$$

Since, $b = 0$

$$\therefore PV = RT - \frac{a}{V} \quad (y = c + MX \text{ form})$$

when PV is plotted against $(1/V)$, we get straight line with negative slope, i.e. $(-a)$.

$$\therefore \text{Slope} = \frac{21.1 - 24.6}{3} = -1.5 = -a$$

$$a = 1.5$$

34. (C)

Work done = Area under BC line

$$\frac{1}{2}(6P_1 + P_1) \times 2V_1 = 7P_1V_1$$

35. (B)

$$G = H - TS$$

$$G = U + PV - TS$$

$$\Delta G = \Delta U + P\Delta V + V\Delta P - T\Delta S - S\Delta T$$

From the first and second laws,

$$T\Delta S = \Delta U + P\Delta V$$

$$\therefore \Delta G = V\Delta P - S\Delta T$$

At constant pressure, $\Delta P = 0$

$$\frac{\Delta G}{\Delta T} = -S$$

From eqns. (i) and (ii)

$$G = H + T \frac{\Delta G}{\Delta T} \text{ or } G = H + T \left(\frac{\partial G}{\partial T} \right)_P$$

$$-\frac{H}{T^2} = -\frac{G}{T^2} + \frac{1}{T} \left(\frac{\partial G}{\partial T} \right)_P = \left[\frac{\partial(G/T)}{\partial T} \right]_P$$

$$H = -T^2 \left[\frac{\partial(G/T)}{\partial T} \right]_P$$

36. (B)

$$\Delta S = 2.303 nR \log_{10} \left(\frac{V_2}{V_1} \right)$$

$$= 2.303 \times 2 \times 8.314 \log_{10} \left(\frac{100}{10} \right)$$

$$= 38.3 \text{ J mol}^{-1} \text{ K}^{-1}$$

37. (A)

$$m\text{-moles of HCl} = 20 \times 0.1 = 2$$

$$m\text{-moles of MgO reacted with HCl} = \frac{2}{2} \Rightarrow 1$$

$$\text{mass of MgO present} = 1 \times 40 \text{ mg}$$

$$\therefore \% \text{ of MgO} = \frac{40}{320} \times 100 = 12.5\%$$

38. (C)

$$\frac{r_{\text{CH}_3\text{OCH}_3}}{r_{\text{CH}_4}} = \sqrt{\frac{M_{\text{CH}_4}}{M_{\text{CH}_3\text{OCH}_3}}} \times \frac{P_{\text{CH}_3\text{OCH}_3}}{P_{\text{CH}_4}}$$

$$= \sqrt{\frac{16}{46}} \times \frac{0.8}{0.2} = 2.36 : 1$$

39. (A)

$$\frac{\partial}{\partial T} \ln K_p = \frac{\Delta H}{RT^2}$$

$$\frac{\partial}{\partial T} \ln K = \frac{E_a}{RT^2}$$

40. (C)

$$0.40 = aT_1^3 + bT_1$$

$$0.40 = a \times (1000) + b \times 10$$

$$0.4 = 1000a + 10b$$

$$0.92 = aT_2^3 + bT_2$$

$$\Rightarrow 0.92 = a \times 8000 + 20b$$

from Eqs. (1) and (2)

$$a = 2 \times 10^{-3}, b = 0.038$$

$$S_m = \int \frac{aT^3 + bT}{T} \cdot dT$$

$$= \frac{a[T_2^3 - T_1^3]}{3} + b[T_2 - T_1]$$

$$= 0.813 \text{ J/K - mol}$$

41. (C)

42. (C)

Compound 'A' can show only optical isomerism due to presence of a chiral carbon, 'B' can show only geometrical isomerism, 'C' can show both geometrical as well as optical isomerism, D can show only optical isomerism.

43. (C)

(A) & (B) Cis-1,3-dimethyl cyclohexane & (D) is trans-1,4-dimethyl cyclohexane

44. (D)

The compound P is identical with (D) because both are meso. (A, B & C) are optically active.

45. (C)

46. (D)

47. (A)

melting point \propto closed packing .


p- substituted benzene has linear geometry and such molecules are closely packed in their solid state.

48. (C)

49. (B)


50. (B)

51. (C)




Chlorotetracyano superoxoferrate (II) anion.

52. (C)




53. (A)



54. (D)

Due to decreasing trend of I.E. reactivity increase along the group but due to standard electrode potential reactivity of halogen in decreasing order along the group..

55. (B)

 $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ exist as $\text{Na}_2 [\text{B}_4\text{O}_5(\text{OH})_4] \cdot 8\text{H}_2\text{O}$

 sp^2 hybridised boron atom = 2 ;

 sp^3 hybridised boron atom = 2.

56. (A)

 $\Delta H_{\text{ion.}} = -\Delta H_{\text{eg}}$

57. (B)

Conceptual.



58. (C)

 $[\text{Zn}(\text{Br})_4]^{2-} = \text{sp}^3$ tetrahedral. Diamagnetic.


59. (B)

Strength of ligands \propto stability of complex.
 $\text{H}_2\text{O} < \text{NH}_3 < \text{NO}_2^-$ is correct order of stability

60. (C)

(C) sp^2 'S' Chr = 33 %

'P' Chr = 66%

(D) Hybridized orbitals always form σ bond due to π bond form by pure orbital only.

MATHEMATICS

61. (A)


Let $\log_3 x = t$, then

$$2t^2 - |t| = -p$$

if $t \geq 0$, $2t^2 - t = -p$

if $t < 0$, $2t^2 + t = -p$

$$\therefore \text{ for 4 distinct roots } -\frac{1}{8} < -p < 0 \Rightarrow 0 < p < \frac{1}{8}$$



62. (C)

$$m_{SP} = \frac{2 \tan \alpha}{\tan^2 \alpha - 1} = -\tan 2\alpha = \tan(\pi - 2\alpha)$$

since $0 < \pi - 2\alpha < \pi$

63. (A) $\sin x \sqrt{8} |\cos x| = 1$

$$2 \sin x |\cos x| = \frac{1}{\sqrt{2}}$$

If $-\frac{\pi}{2} < x < \frac{\pi}{2}$

$$\sin 2x = \sin \frac{\pi}{4} \Rightarrow 2x = n\pi + (-1)^n \frac{\pi}{4} \Rightarrow x = \frac{\pi}{8}, \frac{3\pi}{8} \Rightarrow \text{common difference} = \frac{\pi}{4}$$

64. (B)

$$f(x) = \sqrt{3} \sin x - \cos x + 2 = 2 \sin \left(x - \frac{\pi}{6} \right) + 2$$

Since $f(x)$ is one-one and onto, f is invertible.


$$\text{Now } f \circ f^{-1}(x) = x \Rightarrow 2 \sin \left(f^{-1}(x) - \frac{\pi}{6} \right) + 2 = x$$

$$\Rightarrow \sin \left(f^{-1}(x) - \frac{\pi}{6} \right) = \frac{x}{2} - 1 \Rightarrow f^{-1}(x) = \sin^{-1} \left(\frac{x}{2} - 1 \right) + \frac{\pi}{6}$$

Because $\left| \frac{x}{2} - 1 \right| \leq 1$ for all $x \in [0, 4]$

Hence (B) is the correct answer.

65. (C)

Clearly $g(x)$ is symmetric about $x = 1/2$ 

66. (B)

$$x = 0 \Rightarrow f(2) = 2f(0) - f(1) = 2 \times 2 - 3 = 1$$

$$x = 1 \Rightarrow f(3) = 6 - 1 = 5$$

$$x = 2 \Rightarrow f(4) = 2f(2) - f(3) = 2 \times 1 - 5 = -3$$

$$x = 3 \Rightarrow f(5) = 2f(3) - f(4) = 2(5) - (-3) = 13$$

Hence (B) is correct answer.

67. (B)

$$3 = \lim_{x \rightarrow 0} (1 + a \sin x)^{\csc x} \quad [1^\infty \text{ form}] \Rightarrow \lim_{x \rightarrow 0} e^{\csc x \cdot a \sin x} = e^a$$

$$\therefore e^a = 3 \Rightarrow a = \log_e 3 = \ln 3.$$

68. (B)

$$\text{Using LMVT, } \frac{\tan^{-1} \beta - \tan^{-1} \alpha}{\beta - \alpha} = \frac{1}{1+c^2}$$

where, $0 < \alpha < c < \beta < \sqrt{3}$

$$\text{So, } \frac{1}{4} < \frac{1}{1+c^2} < 1$$

$$\Rightarrow \frac{1}{4} < \frac{\tan^{-1} \beta - \tan^{-1} \alpha}{\beta - \alpha} < 1 \Rightarrow \frac{1}{4} < \frac{\tan^{-1} \left(\frac{\beta - \alpha}{1 + \alpha\beta} \right)}{\beta - \alpha} < 1$$

$$\Rightarrow 1 < \frac{\beta - \alpha}{\cot^{-1} \left(\frac{1 + \alpha\beta}{\beta - \alpha} \right)} < 4$$

69. (C)

Equation of Normal through A is :

$$y = m_1x - 2am_1 - am_1^3$$

for the parabola $y^2 = 4ax$ where $4a = 1071$


$$\text{Here } N \equiv \left(\frac{1071}{2} + \frac{1071}{4}m_1^2, 0 \right)$$

$$\text{If } B = (h, k), \text{ then } k = \frac{1071}{4}m_1$$

$$\text{But since } k = m_1h - 2am_1 - am_1^3$$

$$\Rightarrow m_1m_2m_3 = \frac{-k}{a} = \frac{-\frac{1071}{4}m_1}{\frac{1071}{4}} \Rightarrow m_2m_3 = -1$$

Hence option (C)



70. (D)

Intersection of the ellipse is possible if $a > 1$

$$\Rightarrow b^2 - 5b + 7 > 1 \Rightarrow b^2 - 5b + 6 > 0 \Rightarrow b \in (-\infty, 2) \cup (3, \infty)$$

71. (C)

$$[\sin x] + [2\cos x] = -3$$

Possible if $[\sin x] = -1, [2\cos x] = -2$

$$\Rightarrow -1 \leq \sin x < 0, -2 \leq 2\cos x < -1$$

$$\Rightarrow \pi < x < 2\pi \text{ and } \frac{2\pi}{3} < x < \frac{4\pi}{3} \Rightarrow \pi < x < \frac{4\pi}{3}$$

$$\text{Now, } f(x) = \sin x + \cos x = \sqrt{2} \sin\left(x + \frac{\pi}{4}\right)$$

Clearly the range of $f(x)$ is $[-\sqrt{2}, -1)$


72. (D)

Given equation is

$$16x^2 - 8\pi x + \pi^2 + 16 = |\tan x|$$

$$\text{Let } f(x) = 16x^2 - 8\pi x + \pi^2 + 16$$

$$\text{and } g(x) = |\tan x|$$



$$\Rightarrow f(x) = 16 \left[\left(x - \frac{\pi}{4} \right)^2 + 1 \right]$$

or $y = f(x)$ is an upward parabola with vertex $\left(\frac{\pi}{4}, 16 \right)$

Now, from graph it is clear that $f(x)$ and $g(x)$ intersect at infinitely many points.

73. (B)

$$m_2(x) = \min_{0 \leq t \leq x} (t^2 + (x-t)^2) = \min_{0 \leq t \leq x} 2 \left(\left(t - \frac{x}{2} \right)^2 + \frac{x^2}{4} \right) = \frac{x^2}{2}$$

$$m_3(x) = \min_{0 \leq t \leq x} \left(\frac{t^2}{2} + (x-t)^2 \right) = \min_{0 \leq t \leq x} \frac{3}{2} \left(\left(t - \frac{2}{3}x \right)^2 + \frac{2}{9}x^2 \right) = \frac{x^2}{3}$$

...

$$m_n(x) = \frac{x^2}{n} \therefore m_n \left(\frac{1}{\sqrt{n+1}} \right) = \frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$$

$$\therefore \sum_{n=1}^k m_n \left(\frac{1}{\sqrt{n+1}} \right) = \left(\frac{1}{1} - \frac{1}{2} \right) + \left(\frac{1}{2} - \frac{1}{3} \right) + \dots + \left(\frac{1}{k} - \frac{1}{k+1} \right) = 1 - \frac{1}{k+1} = \frac{k}{k+1}$$

74. (D)


$$ae = \sqrt{\sec^2 \theta - \tan^2 \theta} = 1$$

Distance between foci = 2

75. (B)

$$p_1 p_2 = \frac{a^2 b^2}{a^2 + b^2} = \frac{a^2 \cdot a^2 (e^2 - 1)}{a^2 e^2} = 6$$

$$\frac{2a^2}{3} = 6 \Rightarrow a^2 = 9 \Rightarrow a = 3$$

hence $2a = 6$ 

76. (D)

$$f(1) = -6$$

for maximum at $x = 1$

$$\lim_{x \rightarrow 1^-} f(x) = \tan^{-1} \alpha - 5 < -6$$

$$\tan^{-1} \alpha < -1 \Rightarrow \alpha < -\tan 1$$

77. (B)

$$x = \frac{b(a \cos \beta) - a(b \cos \alpha)}{b - a} \quad y = \frac{b(a \sin \beta) - a(b \sin \alpha)}{b - a}$$

$$\Rightarrow \frac{x}{y} = \frac{\cos \beta - \cos \alpha}{\sin \beta - \sin \alpha}$$

$$\Rightarrow \frac{x}{y} = \frac{2 \sin\left(\frac{\beta + \alpha}{2}\right) \sin\left(\frac{\alpha - \beta}{2}\right)}{2 \cos\left(\frac{\beta + \alpha}{2}\right) \sin\left(\frac{\beta - \alpha}{2}\right)}$$

$$\Rightarrow x \cos\left(\frac{\alpha + \beta}{2}\right) = -y \sin\left(\frac{\alpha + \beta}{2}\right)$$

$$\therefore x \cos\left(\frac{\alpha + \beta}{2}\right) + y \sin\left(\frac{\alpha + \beta}{2}\right) = 0$$

78. (A)

Let $x = r \cos \theta$ and $y = r \sin \theta$, then $x^2 + y^2 = r^2$

$$\text{Now } 3x^2 - 4xy + 2y^2 = 12$$

$$\Rightarrow r^2(3 \cos^2 \theta - 4 \cos \theta \sin \theta + 2 \sin^2 \theta) = 12$$

$$\Rightarrow r^2 = \frac{24}{5 + \cos 2\theta - 4 \sin 2\theta}$$

$$\therefore m = \frac{24}{5 + \sqrt{17}} = 3(5 - \sqrt{17}) \text{ and } n = \frac{24}{5 - \sqrt{17}} = 3(5 + \sqrt{17}).$$

79. (C)

A rectangular hyperbola circumscribing a Δ also passes through its orthocentre

if $\left(ct_i, \frac{c}{t_i} \right)$ where $i = 1, 2, 3$ are the vertices of the Δ therefore orthocentre is

$$\left(\frac{-c}{t_1 t_2 t_3}, -ct_1 t_2 t_3 \right), \text{ where } t_1 t_2 t_3 t_4 = 1. \text{ Hence orthocentre is } \left(-ct_4, \frac{-c}{t_4} \right) = (-x_4, -y_4)$$

80. (C)

$$\begin{aligned} f(x) &= \cos^2 x + \cos^2 2x + \cos^2 3x = 1 + \cos^2 x + \cos^2 2x - \sin^2 3x \\ &= 1 + \cos^2 x + \cos 5x \cdot \cos x = 1 + \cos x (\cos x + \cos 5x) \\ \Rightarrow \cos x \cdot \cos 2x \cdot \cos 3x &= 0 \end{aligned}$$

$$\Rightarrow x = (2n-1)\frac{\pi}{2} \text{ or } x = (2n-1)\frac{\pi}{4} \text{ or } x = (2n-1)\frac{\pi}{6}$$

$$\Rightarrow x = \frac{\pi}{2}, \frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{6}, \frac{5\pi}{6} \Rightarrow \text{no. of values of } x = 5$$

81. (D)


Tangent at 't' passes through (h, k)

$$\therefore h + kt^2 = 2ct \dots\dots\dots(i)$$

$$\& \left(-\frac{1}{t^2} \right) \cdot \frac{k}{x} = -1 \dots\dots\dots(ii)$$

from (i) and (ii)

$$\text{Locus is } (x^2 + y^2)^2 = 4c^2 xy$$



82. (D)

Let $\frac{y}{x} = \alpha$, then

$$\alpha \text{ will be minimum or maximum when } y = \alpha x \dots\dots (1)$$

is a tangent to the ellipse

$$x^2 + xy + 2y^2 - 6x - 10y + 14 = 0 \dots\dots (2)$$


Solving (1) and (2),

$$(1 + \alpha + 2\alpha^2)x^2 - 2(3 + 5\alpha)x + 14 = 0$$

$$D = 0 \Rightarrow 4(3 + 5\alpha)^2 - 56(1 + \alpha + 2\alpha^2) = 0$$

$$\Rightarrow 3\alpha^2 - 16\alpha + 5 = 0 \Rightarrow \alpha = \frac{1}{3} \text{ or } 5$$

$$\therefore m = \frac{1}{3}, M = 5$$



83. (A)

$$\begin{aligned} \text{We have, } \int \frac{1}{4e^{-x} - 9e^{2x}} dx &= \int \frac{e^x}{4 - 9e^{2x}} dx \\ &= \int \frac{e^x}{2^2 - (3e^x)^2} dx = \frac{1}{3} \int \frac{3e^x}{2^2 - (3e^x)^2} dx = \frac{1}{3} \int \frac{1}{2^2 - (3e^x)^2} d(3e^x) \quad [\because d(3e^x) = 3e^x dx] \\ &= \frac{1}{3} \int \frac{dt}{2^2 - t^2}, \text{ where } t = 3e^x \\ &= \frac{1}{3} \times \frac{1}{2 \times 2} \log \left| \frac{2+t}{2-t} \right| + c = \frac{1}{12} \log \left| \frac{2+3e^x}{2-3e^x} \right| + c \end{aligned}$$

84. (C)

radius of the first circle is half of the second circle.

\Rightarrow Triangle is equilateral. \Rightarrow Incentre and circumcentre coincides.

$$\Rightarrow \alpha - 2\beta = 1 \text{ and } \alpha + \beta = 2 \Rightarrow (\alpha, \beta) \equiv \left(\frac{5}{3}, \frac{1}{3} \right)$$

85. (B)

$$\text{The locus is } \frac{x^2}{16} - \frac{y^2}{48} = 1 \Rightarrow e = \sqrt{\frac{16+48}{16}} = 2$$

86. (B)

$$f'(x) = 5^x + 7^x$$

$$f''(x) = 5^x \ln 5 + 7^x \ln 7 > 0$$

i.e $f'(x)$ is \uparrow

$$\text{Maximum slope} = 5^2 + 7^2 = 74$$

87. (D)

$$P : (a \sec \theta, a \tan \theta); N : \left[\frac{a}{2} (\sec \theta + \tan \theta), \frac{a}{2} (\sec \theta + \tan \theta) \right]$$

$$\Rightarrow \frac{4h}{a} = 2 \sec \theta + \tan \theta \text{ \& } \frac{4k}{a} = \sec \theta + 2 \tan \theta \Rightarrow x^2 - y^2 = 3a^2/16$$

88. (C)

$$y^2 - 2y + 1 = 4x^2$$

$$\Rightarrow (y-1)^2 = 4x^2 \Rightarrow y-1 = 2x, -2x$$

89. (B)

$$y - y_i = \left(\frac{dy}{dx} \right)_{(x_i, y_i)} (x - x_i)$$

$$-y_i = \frac{-y_i}{2x_i} (x_{i+1} - x_i)$$

$$x_{i+1} - x_i = 2x_i$$

$$x_{i+1} = 3x_i$$

90. (C)

If 'O' be the circumcentre then

$$OP + OR > PR \geq AD = 10$$

Also a circle through ABCD has radius $5\sqrt{2}$.

